

DO-TH 97/10

May 1997

# Next-to-Leading Order Evolution of Polarized Fragmentation Functions

M. Stratmann

Universität Dortmund, Institut für Physik,  
D-44221 Dortmund, Germany

## Abstract

We present a brief description of the determination of the two-loop spin-dependent time-like splitting functions relevant for the NLO evolution of polarized fragmentation functions. Our calculation based on the analytic continuation of the corresponding space-like results obtained within the light-cone gauge method proposed by Curci, Furmanski, and Petronzio. As an application we present an analysis of polarized  $\Lambda$  production in  $e^+e^-$  and  $ep$  collisions.

*Talk presented at the 5th International Workshop on 'Deep Inelastic Scattering and QCD' (DIS '97), Chicago, IL, April 14-18, 1997.*

# NLO Evolution of Polarized Fragmentation Functions

Marco Stratmann

*Universität Dortmund, Institut für Physik  
D-44221 Dortmund, Germany*

**Abstract.** We present a brief description of the determination of the two-loop spin-dependent time-like splitting functions relevant for the NLO evolution of polarized fragmentation functions. Our calculation based on the analytic continuation of the corresponding space-like results obtained within the light-cone gauge method proposed by Curci, Furmanski, and Petronzio. As an application we present an analysis of polarized  $\Lambda$  production in  $e^+e^-$  and  $ep$  collisions.

## GENERAL FRAMEWORK

In analogy to the familiar space-like (S) structure function  $g_1^{(S)}(x, Q^2)$  appearing in longitudinally polarized DIS one can define a similar time-like (T) structure function  $g_1^{(T)}(z, Q^2)$ , where  $q^2 \equiv Q^2 > 0$ , which describes the helicity transfer in single-particle inclusive  $e^+e^-$  annihilation processes (SIA), e.g.,

$$\frac{d\Delta\sigma^{e^+e^- \rightarrow HX}}{dydz} \equiv \frac{d\sigma^{e^-(+), H(+)} }{dydz} - \frac{d\sigma^{e^-(+), H(-)} }{dydz} = \frac{6\pi\alpha^2}{Q^2}(2y-1)g_1^{(T)}(z, Q^2) . \quad (1)$$

where  $z = 2p_H \cdot q/Q^2$ ,  $y = p_H \cdot p_{e^-}/p_H \cdot q$  and  $(\pm)$  denotes the helicities of the  $e^-$  and  $H$ . In terms of the spin-dependent fragmentation functions  $\Delta D_f^H(z, Q^2) \equiv D_{f(+)}^{H(+)}(z, Q^2) - D_{f(+)}^{H(-)}(z, Q^2)$ , where  $D_{f(+)}^{H(+)}(z, Q^2)$  ( $D_{f(+)}^{H(-)}(z, Q^2)$ ) is the probability for finding a hadron  $H$  with positive (negative) helicity in a parton  $f$  with positive helicity at a mass scale  $Q$ , carrying a fraction  $z$  of the parent parton's momentum, we can write  $g_1^{(T)}$  to NLO as [1] (the symbol  $\otimes$  denotes the usual convolution)

$$g_1^{(T)}(z, Q^2) = \sum_q e_q^2 \left\{ \left[ \Delta D_q^H + \Delta D_{\bar{q}}^H \right] \otimes \Delta \mathcal{C}_q^{(T)} + 2\Delta D_g^H \otimes \Delta \mathcal{C}_g^{(T)} \right\} (z, Q^2) . \quad (2)$$

The  $Q^2$ -evolution of the  $\Delta D_f^H$ , predicted by QCD, is again similar to the space-like case. The singlet evolution equation, e.g., schematically reads

$$\frac{d}{d \ln Q^2} \begin{pmatrix} \Delta D_\Sigma^H \\ \Delta D_g^H \end{pmatrix} (z, Q^2) = \left[ \Delta \hat{P}^{(T)} \otimes \begin{pmatrix} \Delta D_\Sigma^H \\ \Delta D_g^H \end{pmatrix} \right] (z, Q^2) , \quad (3)$$

with  $\Delta D_\Sigma^H \equiv \sum_q (\Delta D_q^H + \Delta D_{\bar{q}}^H)$ . It should be recalled that the off-diagonal entries in the singlet evolution matrices  $\Delta \hat{P}^{(S,T)}$  interchange their role when going from the space- to the time-like case, see, e.g., [2,1].

As a manifestation of the so-called Gribov-Lipatov relation [3] the space- and time-like splitting functions are equal in LO. Furthermore they are related by analytic continuation of the space-like splitting functions (Drell-Levy-Yan relation (ACR) [4]) which can be schematically expressed as ( $z < 1$ )

$$\Delta P_{ij}^{(T)}(z) = z \mathcal{AC} \left[ \Delta P_{ji}^{(S)}(x = \frac{1}{z}) \right], \quad (4)$$

where the operation  $\mathcal{AC}$  analytically continues any function to  $x \rightarrow 1/z > 1$  and correctly adjusts the color factor and the sign depending on the splitting function under consideration [1], e.g.,

$$\Delta P_{qq}^{(T)}(z) = -z \Delta P_{qq}^{(S)}(\frac{1}{z}), \quad \Delta P_{gq}^{(T)}(z) = \frac{C_F}{2T_f} z \Delta P_{qg}^{(S)}(\frac{1}{z}), \quad \dots \quad (5)$$

These LO relations are based on symmetries of tree diagrams under crossing, and should be therefore no longer valid when going to NLO. Fortunately [5], the breakdown of the ACR arising beyond LO is essentially due to kinematics and can be rather straightforwardly detected within the light-cone gauge method used in [2,5–7] to calculate the space-like splitting functions.

## UNDERSTANDING OF THE ACR BREAKING

The general strategy of the light-cone gauge method is based on a rearrangement of the perturbative expansion of a partonic cross section into a finite hard part and a process-*independent* part  $\Gamma_{ij}$  which contains all (and only) mass singularities [8,5]. Using dimensional regularization the  $\overline{\text{MS}}$  evolution kernels appear order by order as the residues of the *single*  $1/\epsilon$  poles in  $\Gamma_{ij}$  [5]. The difference between the space- and the time-like  $\Gamma_{ij}$ -kernels essentially amounts to relative extra phase space factors ( $z^{-2\epsilon}$ ) in the time-like case<sup>1</sup> [5,1]. All this gives on aggregate for  $z < 1$  for the AC of the spin-dependent space-like kernels [1]

$$\Delta \Gamma_{qq}^{(T)}(z, \frac{1}{\epsilon}) = -z^{1-2\epsilon} \Delta \Gamma_{qq}^{(S)}(\frac{1}{z}, \frac{1}{\epsilon}), \quad \Delta \Gamma_{gq}^{(T)}(z, \frac{1}{\epsilon}) = \frac{C_F}{2T_f} z^{1-2\epsilon} \Delta \Gamma_{qg}^{(S)}(\frac{1}{z}, \frac{1}{\epsilon}), \dots \quad (6)$$

Obviously, higher ( $1/\epsilon^2$ ) pole terms in  $\Delta \Gamma_{ij}^{(S)}$  will generate additional contributions to the relevant *single* pole part of  $\Delta \Gamma_{ji}^{(T)}$  if combined with the factors  $z^{-2\epsilon}$  in (6). To extract the contributions that break the ACR one can easily go through the NLO calculation of the  $\Delta \Gamma_{ij}^{(S)}$  [7] graph by graph picking

---

<sup>1)</sup> In the unpolarized case a further difference arises [1] through the necessary adjustment of the different spin-average factors for gluons and quarks in  $d = 4 - 2\epsilon$  dimensions.

up all  $1/\epsilon^2$  pole terms. Apart from the slightly subtle case of the so-called 'ladder-subtraction' topology [7] this is a rather straightforward task [1]. To obtain the desired final result for  $\Delta P_{ij}^{(T)}$  [1] one has to combine the results of the AC of the NLO splitting functions  $\Delta P_{ij}^{(S)}$  [9,7], by using the operation  $\mathcal{AC}[\dots]$  defined above, with the contributions that break the ACR. Finally, the missing endpoint contributions, i.e., the terms  $\sim \delta(1-z)$ , are exactly the same as in the space-like situation [5,2]. In the same way the hard subprocess cross sections  $\Delta C_i^{(T)}$  in (2) can be obtained via analytic continuation (see ref. [1] for details). The final results for the  $\Delta P_{ij}^{(T)}$  and  $\Delta C_i^{(T)}$  are too long to be listed here but can be found in ref. [1]<sup>2</sup>.

The rather simple and transparent structure of the ACR breaking part is a hint that there could be a more straightforward way of linking the time- and the analytically continued space-like NLO quantities. The key observation for such considerations is to notice that although the usual 4-dim. LO splitting functions obey the ACR rule, the rule must break down for their  $d = (4 - 2\epsilon)$ -dimensional counterparts as an immediate consequence of eq.(6) [1]. It turns out [1] that the breakdown of the ACR beyond LO in the  $\overline{\text{MS}}$  scheme is entirely driven by this breaking such that it can be accounted for by a simple factorization scheme transformation [1].

Finally, our results for  $\Delta P_{ij}^{(T)}$  (as well as the corresponding unpol. results given in [5,2]) fulfil the so-called SUSY relation which links all singlet splitting functions in a remarkably simple way in the limit  $C_F = N_C = 2T_f \equiv N$  (cf. [9,7]) if they are properly transformed to a regularization method which respects SUSY such as dimensional reduction [10]. The validity of the SUSY relation [1] serves as an important check for the correctness of our results.


## APPLICATION: POLARIZED $\Lambda$ PRODUCTION

The most likely candidate for a measurement of polarized fragmentation functions is the  $\Lambda$  baryon due to its self-analyzing decay  $\Lambda \rightarrow p\pi^-$ . In [11] a strategy was proposed for extracting the  $\Delta D_f^\Lambda$  in SIA  $e^+e^- \rightarrow \Lambda X$ . At high enough energies, no beam polarization is needed since the parity-violating coupling  $q\bar{q}Z$  automatically generates a net polarization of the quarks. In Fig. 1a we compare the first results for such a measurement of the cross section asymmetry  $A_{SIA}^\Lambda$  on the  $Z$ -pole by ALEPH and DELPHI [12] with two conceivable NLO models for the  $\Delta D_f^\Lambda$  (the corresponding LO results are very similar) [13]. Scenario 1, where  $\Delta D_s^\Lambda(z, \mu^2) = z^\alpha D_s^\Lambda(z, \mu^2)$ ,  $\Delta D_{f \neq s}^\Lambda(z, \mu^2) = 0$ , is based on the naive quark model whereas for scenario 2 we also allow for a non-vanishing  $\Delta D_{u,d}^\Lambda$  input as discussed in [11]. In both cases the inputs for the evolution are specified at some low starting scale  $\mu^2 \simeq \mathcal{O}(0.3 \text{ GeV}^2)$  (for more details see ref. [13] where also a model for the unpolarized  $D_f^\Lambda$  can be found).

---

<sup>2)</sup> We also have recalculated the unpolarized NLO time-like Altarelli-Parisi kernels where we fully agree with the results given in [5,2].

Apart from SIA the possibility of extracting the  $\Delta D_f^\Lambda$  in semi-inclusive DIS (SIDIS), i.e.,  $ep \rightarrow e\Lambda X$ , has also been discussed recently [14]. Here, either a longitudinally polarized lepton beam or a polarized nucleon target would be required. Such kind of measurement appears possible e.g. for the HERMES experiment [15]. In Fig. 1b we present LO and NLO predictions for the cross section asymmetry  $A_{SIDIS}^\Lambda$  in  $\vec{ep} \rightarrow \vec{\Lambda}X$  as a function of  $x$  at  $Q^2 = 3 \text{ GeV}^2$  and where we have integrated over  $0.3 \leq z \leq 0.9$  [13]. Such a measurement by HERMES would allow, in principle, to discriminate between the two proposed scenarios for the polarized  $\Lambda$  fragmentation functions  $\Delta D_f^\Lambda$ .



**FIGURE 1.** a) Comparison of recent data for  $A_{SIA}^\Lambda$  [13] with the results obtained for two conceivable scenarios for the  $\Delta D_f^\Lambda$ ; b) LO and NLO predictions for  $A_{SIDIS}^\Lambda$  in  $\vec{ep} \rightarrow \vec{\Lambda}X$ .

*Acknowledgements:* It is a pleasure to thank W. Vogelsang and D. de Florian for a fruitful collaboration. This work was partly supported by the BMBF Bonn.

## REFERENCES

1. M. Stratmann and W. Vogelsang, hep-ph/9612250, to appear in Nucl. Phys. **B**.
2. W. Furmanski and R. Petronzio, Phys. Lett. **97B** (1980) 437.
3. V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. **15** (1972) 438 and 675.
4. S.D. Drell et al., Phys. Rev. **187** (1969) 2159; Phys. Rev. **D1** (1970) 1617.
5. G. Curci, W. Furmanski, and R. Petronzio, Nucl. Phys. **B175** (1980) 27.
6. R.K. Ellis and W. Vogelsang, hep-ph/9602356.
7. W. Vogelsang, Phys. Rev. **D54** (1996) 2023; Nucl. Phys. **B475** (1996) 47.
8. R.K. Ellis et al., Phys. Lett. **78B** (1978) 281; Nucl. Phys. **B152** (1979) 285.
9. R. Mertig and W.L. van Neerven, Z. Phys. **C70** (1996) 637.
10. W. Siegel, Phys. Lett. **84B** (1979) 193.
11. M. Burkardt and R.L. Jaffe, Phys. Rev. Lett. **70** (1993) 2537.
12. ALEPH collab., Phys. Lett. **B374** (1996) 319; DELPHI collab., DELPHI 95-86.
13. D. de Florian, M. Stratmann, and W. Vogelsang, in preparation.
14. R.L. Jaffe, Phys. Rev. **D54** (1996) 6581; D. de Florian and R. Sassot, Nucl. Phys. **B488** (1997) 367.
15. HERMES collab., DESY-PRC 90/01; M. Düren, DESY-HERMES 95-02.